

Radioactivity

Lecture 8

Biological Effects of Radiation

Studies of impact of ionizing radiation on the human body - Hiroshima -

US-Japanese teams medical tests, autopsies, human organ analysis, on-site radioactivity measurements ...




autopsy




Radiation Exposure Types


Irradiation




External Contamination



Internal Contamination




Schematic Model of Radionuclide Uptake



Deposition depends on chemical characteristics of isotopes

Basic DNA structure


Genetics is the science of heredity and variation in biological systems! It is based on the gradual change of the gene structure in the DNA Deoxyribonucleic acid, which contains the genetic code & instructions for the biological system in the sequence of the four A, S, G, T types of nucleotides




The DNA forms a double stranded molecule, the double helix. Pair coupling of two nucleotides on the opposite strand :


A with T,
G with C.

Humans have $\sim 3 \cdot 10^9$ base pairs



Radiation interacting with cell molecules






X & Y chromosome

Three types of DNA ...

Nuclear DNA (blue print of body structure...)
inherited by cell division from parents


Y chromosome DNA (determines the male sex)
inherited from father to son
(male XY chromosome, female XX chromosome)

Mitochondrial DNA (outside the cell nucleus)
inherited from mother to children



Damage could cause cancer of
body organ or impact genetic
information for next generation,

⇒ **mutation!!!**




U.S. National Library of Medicine



U.S. National Library of Medicine

Energy dependence of radiation damage



Linear energy transfer (LET): amount of energy deposited per unit track length


Sequence of Events in Indirect Action

$T_{1/2}$ in sec

10^{-15}

10^{-5}

10^{-5}



Incident X-ray photons



Fast electrons



Ion radicals



Free radicals





Macromolecular changes from
breakage of chemical bonds



Possible biological effects

days
generation
years

- cell killing
- mutation
- carcinogenesis



What are the high level radiation effects?

RADIATION EFFECTS

Measurements in millisieverts (mSv). Exposure is cumulative.

Potentially fatal radiation sickness. Much higher risk of cancer later in life.

10,000 mSv: Fatal within days.

5,000 mSv: Would kill half of those exposed within one month.

2,000 mSv: Acute radiation sickness.

No immediate symptoms. Increased risk of serious illness later in life.

1,000 mSv: 5% higher chance of cancer.

400 mSv: Highest hourly radiation recorded at Fukushima. Four hour exposure would cause radiation sickness.

100 mSv: Level at which higher risk of cancer is first noticeable

No symptoms. No detectable increased risk of cancer.

20 mSv: Yearly limit for nuclear workers.


10 mSv: Average dose from a full body CT scan

9 mSv: Yearly dose for airline crews.

3 mSv: Single mammogram

2 mSv: Average yearly background radiation dose

0.1 mSv: Single chest x-ray



EYES High doses can trigger cataracts months later.

THYROID Hormone glands vulnerable to cancer. Radioactive iodine builds up in thyroid. Children most at risk.

LUNGS Vulnerable to DNA damage when radioactive material is breathed in.

STOMACH Vulnerable if radioactive material is swallowed.

REPRODUCTIVE ORGANS High doses can cause sterility.

SKIN High doses cause redness and burning.

BONE MARROW Produces red and white blood cells. Radiation can lead to leukaemia and other immune system diseases.

Acute Radiation Syndrome Recognition

- Signs and symptoms experienced by individuals exposed to acute whole body irradiation.
- Data collected largely through Japanese atomic bomb survivors at Hiroshima and Nagasaki.
- Limited number of accidents at nuclear installations.
- Clinical radiotherapy.
- Well-characterized animal data base.
- Lethal Dose is $\geq 8 \text{ Gy} = 800 \text{ rad}$.
- LD_{50} dose of human is $\sim 4.5 \text{ Gy} = 450 \text{ rad}$.





Early Lethal Effects

Hematopoietic syndrome:

- Hair loss (Epilation) and radiation sickness >2 Gy
- Purpura, or bleeding under the skin.
- Suppresses normal bone marrow and spleen functions.
- Lens opacity, blindness
- Symptoms associated with hematopoietic syndrome are: chill, fatigue, hemorrhages, ulceration, infection and anemia. Death at $D > 8$ Gy unless receive bone marrow transplant.



Gastrointestinal syndrome:

- Occurs at dose >10 Gy of gamma-rays or its equivalence.
- Death usually occurs within 3 to 10 days.
- Symptoms due largely to depopulation of the epithelial lining of the GI tract by radiation.
- No human has survived radiation dose >10 Gy.
- Clinical symptoms include nausea, vomiting, and prolonged diarrhea, dehydration, loss of weight, complete exhaustion, and eventually death.



Human lethality as function of dose

A 50% lethality is reached with an accumulated dose of 450 cGy = 450 rad = 4.5 Gy. A dose of 100 rad (1 Sv) is survivable. A 800 rad (8 Gy) dose is lethal.



Alexander Litvinenko

Litvinenko died on radiation poisoning by a small amount of ^{210}Po after 22 days. ^{210}Po is an alpha emitter that has a half-life of 138.4 days; it decays directly to its stable daughter isotope, ^{206}Pb . The alpha particles have an energy of 5.307 MeV. Calculate the minimum amount of ^{210}Po Litvinenko must have eaten assuming a deadly exposure to 8 Gy.

(1 MeV = $1.6 \cdot 10^{-13}$ Joule)

(1 Gy = 1 J/kg)



$$A = \lambda \cdot N = \frac{\ln 2}{T_{1/2}} \cdot N = 5 \cdot 10^{-3} \cdot N [d^{-1}]$$

$$210 \text{ g} = 6.022 \cdot 10^{23} \text{ particles} \quad N = m[\text{g}] \cdot 2.87 \cdot 10^{21} \text{ particles}$$

$$D = \frac{E}{m} \cdot A \cdot t = \frac{5.307 \text{ MeV}}{80 \text{ kg}} \cdot A \cdot t = \frac{4.25 \cdot 10^{-15} \text{ J}}{80 \text{ kg}} \cdot N \cdot t \quad t = 22 \text{ d}$$

$$D = 1.17 \cdot 10^{-15} \text{ Gy} \cdot N = 3.35 \cdot 10^6 \cdot m[\text{g}] \text{ Gy}$$

$$m[\text{g}] = \frac{8 \text{ Gy}}{3.35 \cdot 10^6} = 2.4 \cdot 10^{-6} \text{ g} \quad A = 3.43 \cdot 10^{13} \text{ d}^{-1} = 4 \cdot 10^9 \text{ s}^{-1} = 4 \cdot 10^9 \text{ Bq}$$

Yasser Arafat

Measurements suggest small amounts of about 2 pico-gram ($1 \text{ pg} = 10^{-12} \text{g}$) of ^{210}Po in the body of Yasser Arafat. Our natural amount of ^{210}Po corresponds to 60 Bq. **By how much did Yasser Arafat exceed this limit? What was his dose?**

$$N = \frac{2 \cdot 10^{-12}}{60} \cdot 6.022 \cdot 10^{23} = 2 \cdot 10^{10} \text{ } ^{210}\text{Po atoms}$$

$$A = \lambda \cdot N = \frac{\ln 2}{T_{1/2}} \cdot N = \frac{0.69}{138.4 \cdot 24 \cdot 3600} \cdot 2 \cdot 10^{10} \text{ s}^{-1}$$

$$A \approx 1170 \text{ Bq} \quad D = 5.3 \cdot 10^{-17} \text{ Gy} \cdot A \cdot t = 6.2 \cdot 10^{-14} \cdot t$$

$$t = 2 \text{ y} = 6.3 \cdot 10^7 \text{ s} \quad D \approx 4 \mu\text{Gy}$$

About a factor of 20 higher than the natural ^{210}Po activity of the human body! High uncertainties in these data because of the two year time period between death and exhumation. The detected amount is only 3% of the original ^{210}Po amount.




Improving forensic investigation for polonium poisoning.
Froidevaux P, et al. Lancet. 2013 Oct 12;382(9900):1308.

$$\frac{N}{N_0} = e^{-\lambda t} = e^{-\frac{\ln 2}{138.4} \cdot 712} = 0.03$$

$$H = \frac{4 \cdot 10^{-6}}{0.03} * 20 = 2.67 \text{ mSv}$$

What are the low level radiation effects?



LNT Definition

- The LINEAR NO-THRESHOLD MODEL (LNT) is a model used in radiation protection to estimate the long-term, biological damage caused by ionizing radiation
- It is assumed that the damage is directly proportional (“linear”) to the dose of radiation, at all dose levels
- Radiation is always considered harmful with no safety threshold

Fierce arguments due to lack of data and statistical interpretation!


Pro: BEIR and UNSCEAR

Against: medical and radiological experts



Hormesis or Linear No Threshold model

The LNT model is the recommended model for extrapolating cancer probability from known high dose exposures to unknown low-dose regions of exposure. It is the recommended approach by the Atomic Energy Commission. This approach is questioned by some radiation experts who claim that low level radiation may have beneficial effects, stimulating the activation of repair mechanisms that protect against disease. This phenomenon is observed for many poisonous substances. A number of compensatory and reparatory mechanisms are activated in response to the damage caused by ionizing radiation. These include up-regulation of antioxidant responses, scavenging of damaged cells that may undergo tumorigenesis, activation of enzymatic DNA repair mechanisms, and activation of the immune system to help recognize and transform mutated cells.

Possible hormesis effect below 0.2 Sv?




Increased risk of solid cancer as function of dose in atomic bomb survivors.



Incidence of leukemia as function of dose in atomic bomb survivors in Japan.

Hormesis defenders



Harold Agnew (1921-2013)



Edward Teller (1908-2003)




Miskolc Tapolca, Hungary



Bad Gastein, Austria

Enzymatic Repair Mechanisms




The capability of repair depends on the kind of radiation, the extent of damage, the capability of enzymes to identify and chemically neutralize aggressive radicals (OH^-). Also important is the cycle time for cell division. In fast cycles less time for repair. Fast cycling blood cells are more prone to radiation damage, but fast growing cancer cells are more vulnerable to radiation bombardment.

The repair mechanism damaged sequence from helix strand. In case of newly synthesized, based on secondary information carriers. If those are not around, the repair is made on enzyme guesswork. This can cause changes in information work, causing mutation. Unfavorable mutations are naturally eliminated. Favorable ones lead to evolution!





by enzymes is based on copying the the identical sequence in the second larger damage, whole sections are

DNA Division and Copying



DNA divisions happen continually with cell divisions. Each of the new cells must contain identical DNA structure and information. The DNA division produces replica of the original DNA molecule using each of the strands as a template! The frequency of the cell division depends on cell. Blood cells have high replication rates, so have cancer cells! Changes in DNA information happens during the division causing a mutation of the cell information. This mutation can be spontaneous or induced depending on the cause for the error in replication!

Mutation



A mutation is any alteration of a genome. It includes substitutions, insertions, and deletions causing a change in the information transfer during cell division. There are 23 pairs of chromosomes in the human genome. There are a total of six billion places where a mutation can happen.





Natural mutation rate is characterized as balance between spontaneous mutation and natural selection and is about $2 \cdot 10^{-10}$ Mutationen per generation and base. With $3 \cdot 10^9$ bases per genom one anticipates ~ 5 mutations per genom and generation. With about 400 cell divisions generations for an adult male we get 2000 mutations. The natural mutation rate is spontaneous, while a radiation induced mutation is deterministic and increases linearly towards higher doses. **But how is the dose dependence at low dose?**

Long range genetic effects




Chromosomes observed during cell division. Abnormal ones are marked by grey arrow.



Observed increase with dose indicates long term genetic effects

Is there a natural threshold, e.g. where random mutation mechanisms are exceeded by radiation induced mutation?

Internal radiation exposure is at 30 mSv/year (mainly γ rays). Life time dose is 2.1 Sv (Gy)




Radiation-induced Mutagenesis


- Radiation *DOES NOT* produce new, unique mutations, but increases the incidence of the same mutations that occur spontaneously.
- Mutation incidence in humans is *DOSE* and *DOSE-RATE* dependent.
- A dose of 1 rem (10 mSv) per generation increases background mutation rate by 1%.
- Information on the genetic effects of radiation comes almost entirely from animal and *IN VITRO* studies.
- Children of A-bomb survivors from Hiroshima and Nagasaki fail to show any significant genetic effects of radiation.
- Other radiation exposed workers (miners, air personnel, radiation workers) also don't show any statistical significant radiation induced health effects (cancer rate).

Epidemiological studies on large population groups

Impact of Radon level on cancer




Distribution of radon emission in the US
(<http://www.kgs.ku.edu/Publications/PIC/pic25.html>).



Cancer rate in US per 100,000 people
according to National Cancer Institute
(<http://statecancerprofiles.cancer.gov/>)



Not a particular convincing correlation, despite corrections for smoker effect.

Cancer risk of Uranium miners



58987 miners recorded from 1946-1990. By 2003 36% had died (20920):
 3016 lung cancer,
 3355 other cancer,
 5141 heart problems
 1742 stroke.

In comparison with death cause distribution in normal population a risk assessment was made.



The excess relative risk per 100 WLM for miners

Stay away from intense radiation sources

ATOMIC

RADIATION hazard

 **TOUCHING or REMOVING
SCRAP OBJECTS IS
PROHIBITED** 

**THIS INCLUDES BLASTED DEBRIS, FUSED SILICA
METAL FRAGMENTS, etc. LOCATED North of *this* POINT**